

## DETECTION OF CHANGES IN LINEAR SEQUENCES

LAJOS HORVÁTH\*

*Department of Mathematics, University of Utah, Salt Lake City, UT 84112, U.S.A.*

(Received May 15, 1995; revised May 1, 1996)

**Abstract.** We discuss the asymptotic properties of some tests to detect possible changes in the mean of linear processes.

*Key words and phrases:* Change-point, linear sequence, dependent observations.

### 1. Introduction and results

Testing for a change in the mean of a sequence of observations is one of the most basic and important problems in change-point analysis. Assuming that the observations are independent normal random variables, Sen and Srivastava (1975*a*, 1975*b*), Worsley (1979, 1986) derived several methods to test for a shift in the normal mean. Yao and Davis (1986), Haccou *et al.* (1988), Gombay and Horváth (1990, 1994) and Horváth (1993*b*) derived the asymptotic distribution of the maximally selected likelihood ratio when changes in the parameters of the observations are tested for. Without assuming any parametric form of the distribution functions of the observations, Page (1954, 1955) and Csörgő and Horváth (1988) studied the detection of changes in the mean. Picard (1985), Kulpelger (1985), Giraitis and Leipus (1990, 1992), Tang and MacNeill (1993), Horváth (1993*a*) and Davis *et al.* (1995) considered tests when the observations are dependent. For the estimation of the time of change in the mean of dependent observations we refer to Picard (1985), Bai (1994) and Horváth and Kokoszka (1995).

We assume that the observations  $\{X_i, 1 \leq i \leq n\}$  satisfy the model

$$X_i = \mu_i + e_i, \quad 1 \leq i \leq n,$$

where the errors  $\{e_i, 1 \leq i \leq n\}$  are given by the linear process

$$(1.1) \quad e_i = \sum_{0 \leq j < \infty} a_j \epsilon_{i-j}.$$

We wish to test the null hypothesis

$$H_0 : \mu_1 = \mu_2 = \cdots = \mu_n$$

---

\* Partially supported by an NSF-NATO East Europe Outreach grant.

against the alternative  $H_A$ : there is an integer  $k^*$ ,  $1 \leq k^* < n$ , such that

$$\mu_1 = \cdots = \mu_{k^*}^* \neq \mu_{k^*+1} = \cdots = \mu_n.$$

The tests for  $H_0$  against  $H_A$  will be based on functionals of

$$Z_n^0(t) = Z_n(t) - tZ_n(1)$$

(cf. Theorems 1.2 and 1.3 for the distribution of weighted supremum functionals), where

$$Z_n(t) = \begin{cases} n^{-1/2} \sum_{1 \leq i \leq (n+1)t} X_i, & \text{if } 0 \leq t < 1 \\ n^{-1/2} \sum_{1 \leq i \leq n} X_i, & \text{if } t = 1. \end{cases}$$

Throughout this paper we assume that

$$(1.2) \quad \{\epsilon_i, -\infty < i < \infty\} \text{ are independent, identically distributed random variables with } E\epsilon_i = 0, \\ 0 < \sigma^2 = E\epsilon_i^2 < \infty \text{ and } E|\epsilon_i|^\nu < \infty \text{ with some } \nu > 2.$$

The random variables  $\epsilon_i$  are smooth with density function  $f$  satisfying

$$(1.3) \quad \sup_{-\infty < s < \infty} \frac{1}{|s|} \int_{-\infty}^{\infty} |f(t+s) - f(t)| dt < \infty.$$

Also, the condition

$$(1.4) \quad a_k = O(k^{-\beta}), \quad \text{as } k \rightarrow \infty, \quad \text{with some } \beta > 3/2$$

holds. Let

$$g(z) = \sum_{0 \leq k < \infty} a_k z^k, \quad z \in \mathbb{C}$$

and assume

$$(1.5) \quad g(z) \neq 0 \quad \text{for all } |z| \leq 1.$$

We note that conditions (1.2) and (1.4) imply that

$$Ee_i = 0 \quad \text{and} \quad \text{var } e_i = \sigma^2 \sum_{0 \leq k < \infty} a_k^2 < \infty.$$

First we obtain the necessary and sufficient condition for the weak convergence of  $Z_n^0$  in weighted metrics. Using weight functions we can improve the power of the tests if the change occurs at the beginning or at the end of the observations (cf. Picard (1985) and Csörgő and Horváth (1988)). Let

$$(1.6) \quad \tau^2 = \sigma^2 \left( \sum_{0 \leq j < \infty} a_j \right)^2 > 0$$

and

$$(1.7) \quad Q = \left\{ q : \inf_{\epsilon \leq t \leq 1-\epsilon} q(t) > 0, q \text{ is non-decreasing in a neighbourhood of zero and non-increasing in a neighbourhood of one} \right\}.$$

The condition for the weak convergence of  $Z_n^0$  in weighted metrics will be given by the integral

$$(1.8) \quad I_{0,1}(q, c) = \int_0^1 \frac{1}{t(1-t)} \exp\left(-\frac{cq^2(t)}{t(1-t)}\right) dt.$$

**THEOREM 1.1.** *We assume that  $H_0$ , (1.1)–(1.6) hold and  $q \in Q$ . If  $I_{0,1}(q, c) < \infty$ , then we can define a sequence of Brownian bridges  $\{B_n(t), 0 \leq t \leq 1\}$  such that*

$$(1.9) \quad \sup_{0 < t < 1} |Z_n^0(t) - \tau B_n(t)|/q(t) = o_P(1).$$

*If  $\limsup_{t \rightarrow 0} t/q(t) < \infty$ ,  $\limsup_{t \rightarrow 1} (1-t)/q(t) < \infty$  and (1.9) holds with a sequence of Brownian bridges, then  $I_{0,1}(q, c) < \infty$  for all  $c > 0$ .*

If we are interested only in the convergence in distribution of weighted supremum functional of  $Z_n^0$ , it can be established under weaker conditions than in Theorem 1.1. Let  $\{B(t), 0 \leq t \leq 1\}$  denote a Brownian bridge.

**THEOREM 1.2.** *We assume that  $H_0$ , (1.1)–(1.6) hold and  $q \in Q$ . Then*

$$(1.10) \quad \sup_{0 < t < 1} |Z_n^0(t)|/q(t) \xrightarrow{D} \tau \sup_{0 < t < 1} |B(t)|/q(t)$$

and

$$(1.11) \quad \sup_{0 < t < 1} Z_n^0(t)/q(t) \xrightarrow{D} \tau \sup_{0 < t < 1} B(t)/q(t)$$

*if and only if  $I_{0,1}(q, c) < \infty$  for some  $c > 0$ .*

The variance of  $Z_n^0(t)$  is proportional to  $\tau^2 t(1-t)$ . However,  $I_{0,1}((t(1-t))^{1/2}, c) = \infty$  for all  $c > 0$ , so the results in Theorems 1.1 and 1.2 cannot be used to get the asymptotic distribution of the standardized statistic

$$(1.12) \quad T_n = \frac{1}{\tau} \sup_{0 < t < 1} |Z_n^0(t)|/(t(1-t))^{1/2}.$$

Bai (1994) pointed out that  $T_n$  is related to the generalized likelihood ratio, if  $\{\epsilon_i, -\infty < i < \infty\}$  are independent, identically distributed normal random variables. Let  $a(t) = (2 \log t)^{1/2}$  and  $b(t) = 2 \log t + \frac{1}{2} \log \log t - \frac{1}{2} \log \pi$ .

THEOREM 1.3. *If  $H_0$  and (1.1)–(1.6) hold, then*

$$\lim_{n \rightarrow \infty} P\{a(\log n)T_n \leq t + b(\log n)\} = \exp(-2e^{-t})$$

*for all  $t$ .*

Theorems 1.1–1.2 will follow from a general invariance principle for tied down partial sums of stationary sequences. The Appendix contains the results and the proofs on weighted convergence of tied down sums of stationary sequence. The proofs of Theorems 1.1–1.3 are given in the next section.

## 2. Proofs of Theorems 1.1–1.3

The first lemma will show that  $\{e_i = \sum_{0 \leq j < \infty} a_j \epsilon_{i-j}, -\infty < i < \infty\}$  is a stationary strongly mixing sequence. For the definition and properties of strongly mixing sequences we refer to Ibragimov and Linnik (1971) and Philipp and Stout (1975).

LEMMA 2.1. *If (1.1)–(1.6) hold, then  $\{e_i, -\infty < i < \infty\}$  is a stationary strongly mixing sequence with mixing coefficient  $\rho(k) = k^{-\alpha}$  with some  $\alpha > 0$ .*

PROOF. The result follows from Gorodetskii (1977) and Withers (1981).

Combining Lemma 2.1 and the strong approximation of partial sums of strongly mixing random variables, we get the next lemma.

LEMMA 2.2. *If (1.1)–(1.6) hold, then we can define a Wiener process  $\{W(t), 0 \leq t < \infty\}$  such that*

$$(2.1) \quad \left| \sum_{1 \leq i \leq k} e_i - \tau W(k) \right| \stackrel{a.s.}{=} o(k^{1/2-\beta})$$

*with some  $\beta > 0$ .*

PROOF. It follows from Lemma 2.1 and Theorem 4 of Kuelbs and Philipp (1980).

Now it is very easy to prove Theorems 1.1 and 1.2.

PROOFS OF THEOREMS 1.1 and 1.2. It follows from Lemma 2.1 that

$$(2.2) \quad n^{-1/2} \sum_{1 \leq i \leq (n+1)t} e_i \xrightarrow{\mathcal{D}[0,1]} \tau W(t)$$

and by the modulus of continuity of Wiener processes (cf. Csörgő and Révész (1981)) we have

$$(2.3) \quad \max_{1 \leq x \leq n} \left| \sum_{1 \leq i \leq x} e_i - \tau W(x) \right| / x^{1/2} = O_P(1).$$

Hence by Lemma 2.2 condition C.2 in the Appendix is also satisfied and therefore Theorems 1.1 and 1.2 follow from Theorems 3.1 and 3.2.

PROOF OF THEOREM 1.3. Using (2.1) we get that

$$(2.4) \quad |Z_n(1)| = O_P(1),$$

and therefore the strong approximation in Lemma 2.2 and Darling and Erdős (1956) yield

$$(2.5) \quad (2\tau^2 \log \log n)^{-1/2} \sup_{0 < t \leq 1/2} |Z_n^0(t)| / (t(1-t))^{1/2} \xrightarrow{D} 1$$

$$(2.6) \quad (2\tau^2 \log \log n)^{-1/2} \sup_{0 < t < 1/\log n} |Z_n^0(t)| / (t(1-t))^{1/2} \xrightarrow{D} 1$$

and

$$(2.7) \quad \sup_{1/\log n \leq t \leq 1/2} |Z_n^0(t)| / (t(1-t))^{1/2} = O_P((\log \log \log n)^{1/2}).$$

By the stationarity of  $\{e_i, 1 \leq i < \infty\}$  we get from (2.5)–(2.7) that

$$(2.8) \quad (2\tau^2 \log \log n)^{-1/2} \sup_{1/2 \leq t < 1} |Z_n^0(t)| / (t(1-t))^{1/2} \xrightarrow{D} 1,$$

$$(2.9) \quad (2\tau^2 \log \log n)^{-1/2} \sup_{1-1/\log n \leq t < 1} |Z_n^0(t)| / (t(1-t))^{1/2} \xrightarrow{D} 1$$

and

$$(2.10) \quad \sup_{1/2 \leq t \leq 1-1/\log n} |Z_n^0(t)| / (t(1-t))^{1/2} = O_P((\log \log \log n)^{1/2}).$$

It follows from (2.4), (2.6) and (2.9) that

$$(2.11) \quad \begin{aligned} & \sup_{0 < t \leq 1/\log n} |Z_n^0(t)| / (t(1-t))^{1/2} \\ &= \max_{1 \leq k \leq n/\log n} k^{-1/2} \left| \sum_{1 \leq i \leq k} e_i \right| + O_P((\log \log n)^{1/2} / \log n) \end{aligned}$$

and

$$(2.12) \quad \begin{aligned} & \sup_{1-1/\log n < t < 1} |Z_n^0(t)| / (t(1-t))^{1/2} \\ &= \max_{n-n/\log n \leq k < n} \frac{1}{(n-k)^{1/2}} \left| \sum_{k < i \leq n} e_i \right| \\ &+ O_P((\log \log n)^{1/2} / \log n). \end{aligned}$$

In light of (2.5)–(2.12), Theorem 1.3 is proven if we show that

$$(2.13) \quad \lim_{n \rightarrow \infty} P \left\{ a(\log n) \frac{1}{\tau} \max_{1 \leq k \leq n/\log n} \frac{1}{k^{1/2}} \left| \sum_{1 \leq i \leq k} e_i \right| \leq t + b(\log n), \right. \\ \left. a(\log n) \frac{1}{\tau} \max_{n-n/\log n \leq k < n} \frac{1}{(n-k)^{1/2}} \left| \sum_{k < i \leq n} e_i \right| \leq s + b(\log n) \right\} \\ = \exp(-(e^{-t} + e^{-s})).$$

Since Lemma 2.2 and Darling and Erdős (1956) yield that

$$(2.14) \quad \lim_{n \rightarrow \infty} P \left\{ a(\log n) \frac{1}{\tau} \max_{1 \leq k \leq n/\log n} \frac{1}{k^{1/2}} \left| \sum_{1 \leq i \leq k} e_i \right| \leq t + b(\log n) \right\} \\ = \exp(-e^{-t})$$

and

$$(2.15) \quad \lim_{n \rightarrow \infty} P \left\{ a(\log n) \frac{1}{\tau} \max_{n-n/\log n \leq k < n} \frac{1}{(n-k)^{1/2}} \left| \sum_{k < i \leq n} e_i \right| \leq s + b(\log n) \right\} \\ = \exp(-e^{-s}),$$

(2.13) is established if we can prove asymptotic independence.

Let

$$X_{i,n} = \sum_{0 \leq j \leq n-3n/\log n} a_j \epsilon_{i-j}, \quad n - n/\log n \leq i \leq n.$$

It is easy to see that  $\{X_i, 1 \leq i \leq n/\log n\}$  and  $\{X_{i,n}, n - n/\log n \leq i \leq n\}$  are independent random vectors. Next we note that

$$X_i - X_{i,n} = \sum_{0 \leq j < \infty} \tilde{a}_j \epsilon_{i-j}, \quad n - n/\log n \leq i \leq n$$

where

$$\tilde{a}_j = \begin{cases} 0, & \text{if } 0 \leq j \leq n - 3n/\log n \\ a_j, & \text{if } n - 3n/\log n < j < \infty. \end{cases}$$

Using the proof of Proposition 1 of Bai (1994) we get

$$(2.16) \quad P \left\{ \max_{n-n/\log n \leq k < n} \frac{1}{(n-k)^{1/2}} \left| \sum_{k < i \leq n} (X_i - X_{i,n}) \right| > x \right\} \\ \leq \frac{9\sigma^2}{x^2} \left( \sum_{0 \leq j < \infty} \tilde{a}_j \right)^2 \left( 1 + \sum_{1 \leq j \leq n/\log n} \frac{1}{j} \right) \\ + \frac{9\sigma^2}{x^2} \sum_{0 \leq j < \infty} \left( \sum_{j < i < \infty} \tilde{a}_j \right)^2 \left( \frac{\log n}{n} + \sum_{1 \leq j \leq n} \frac{1}{j} \right).$$

Applying (1.4) we can find two constants  $C_1$  and  $C_2$  depending on  $\beta$  only, such that

$$(2.17) \quad \left( \sum_{0 \leq j < \infty} \tilde{a}_j \right)^2 \leq C_1 n^{2-2\beta}$$

and

$$(2.18) \quad \sum_{0 \leq j < \infty} \left( \sum_{j < i < \infty} \tilde{a}_j \right)^2 \leq C_2 n^{3-2\beta}.$$

Choosing  $x = 1/\log n$  in (2.16), we get from (1.4), (2.17) and (2.18) that

$$\begin{aligned} & \max_{n-n/\log n \leq k < n} \frac{1}{(n-k)^{1/2}} \left| \sum_{k < i \leq n} X_i \right| \\ &= \max_{n-n/\log n \leq k < n} \frac{1}{(n-k)^{1/2}} \left| \sum_{k < i \leq n} X_{i,n} \right| + O_P(1/\log n), \end{aligned}$$

which gives the asymptotic independence in (2.13).  $\square$

#### Appendix

Let  $\{\xi_k, -\infty < k < \infty\}$  be a sequence of stationary random variables and define

$$S_n(t) = \begin{cases} n^{-1/2} \sum_{1 \leq i \leq (n+1)t} \xi_i, & 0 \leq t < 1 \\ n^{-1/2} \sum_{1 \leq i \leq n} \xi_i, & \text{if } t = 1. \end{cases}$$

We assume that

C.1 there is  $\sigma > 0$  such that

$$S_n(t) \xrightarrow{\mathcal{D}[0,1]} \sigma W(t),$$

where  $\{W(t), 0 \leq t \leq 1\}$  is a Wiener process and

C.2 there are two sequences of Wiener processes  $\{W_n^*(t), 0 \leq t < \infty\}$ ,  $\{W_n^{**}(t), 0 \leq t < \infty\}$  and a sequence of positive numbers  $\sigma_n$  such that

$$\begin{aligned} & \max_{1 \leq x \leq n} \left| \sum_{1 \leq i \leq x} \xi_i - \sigma_n W_n^*(x) \right| / x^{1/2} = O_P(1), \\ & \max_{1 \leq x \leq n} \left| \sum_{-x \leq i \leq -1} \xi_i - \sigma_n W_n^{**}(x) \right| / x^{1/2} = O_P(1). \end{aligned}$$

and

$$0 < \liminf_{n \rightarrow \infty} \sigma_n \leq \limsup_{n \rightarrow \infty} \sigma_n < \infty.$$

We are interested in the weighted approximations of the tied-down partial sums process

$$S_n^0(t) = S_n(t) - tS_n(1), \quad 0 \leq t \leq 1.$$

We find the necessary and sufficient condition for the weak convergence of  $S_n^0(t)$  in weighted metrics, if conditions C.1 and C.2 are satisfied. We recall that the set  $Q$  and the integral  $I_{0,1}(q, c)$  are defined in (1.7) and (1.8), respectively.

**THEOREM A.1.** *We assume that C.1, C.2 hold and  $q \in Q$ . If  $I_{0,1}(q, c) < \infty$  for all  $c > 0$ , then we can define a sequence of Brownian bridges  $\{B_n(t), 0 \leq t \leq 1\}$  such that*

$$(A.1) \quad \sup_{0 < t < 1} |S_n^0(t) - \sigma B_n(t)|/q(t) = o_P(1).$$

*If  $\limsup_{t \rightarrow 0} t/q(t) < \infty$ ,  $\limsup_{t \rightarrow 1} (1-t)/q(t) < \infty$  and (3.1) holds with a sequence of Brownian bridges, then  $I_{0,1}(q, c) < \infty$  for all  $c > 0$ .*

**PROOF.** It follows from C.1 that there is a sequence of Wiener processes  $\{W_n(t), 0 \leq t \leq 1\}$  such that

$$\sup_{0 \leq t \leq 1} |S_n(t) - \sigma W_n(t)| = o_P(1)$$

(cf. Remark 2 in Shorack and Wellner ((1986), p. 49)). Hence

$$(A.2) \quad \sup_{0 \leq t \leq 1} |S_n^0(t) - \sigma B_n(t)| = o_P(1),$$

where

$$(A.3) \quad B_n(t) = W_n(t) - tW_n(1).$$

We assume that  $I_{0,1}(q, c) < \infty$ . Then we have

$$(A.4) \quad \lim_{t \rightarrow 0} q(t)/t^{1/2} = \infty \quad \text{and} \quad \lim_{t \rightarrow 1} q(t)/(1-t)^{1/2} = \infty$$

(cf. Csörgő and Horváth (1993), p. 180). If  $\{B(t), 0 \leq t \leq 1\}$  stands for Brownian bridge, then

$$(A.5) \quad \lim_{\epsilon \rightarrow 0} \sup_{0 < t \leq \epsilon} |B(t)|/q(t) = 0 \quad \text{a.s.}$$

(cf. Csörgő and Horváth (1993), p. 189). Next we show that

$$(A.6) \quad \lim_{\epsilon \rightarrow 0} \limsup_{n \rightarrow \infty} P \left\{ \sup_{0 \leq t \leq \epsilon} |S_n^0(t)|/q(t) > \delta \right\} = 0$$

for all  $\delta > 0$ . It follows from the definition of  $S_n^0(t)$  that

$$\sup_{0 < t \leq \epsilon} |S_n^0(t)|/q(t) \leq \sup_{0 < t \leq \epsilon} |S_n(t)|/q(t) + \sup_{0 < t \leq \epsilon} |S_n(1)t|/q(t)$$

and

$$\sup_{0 < t \leq \epsilon} |S_n(1)t|/q(t) \leq \epsilon^{1/2} |S_n(1)| \sup_{0 < t \leq \epsilon} t^{1/2}/q(t).$$

Using C.1 and (A.4) we obtain that

$$(A.7) \quad \lim_{\epsilon \rightarrow 0} \limsup_{n \rightarrow \infty} P \left\{ \sup_{0 < t \leq \epsilon} |S_n(1)t|/q(t) > \delta \right\} = 0$$

for all  $\delta > 0$ . Next we note that

$$(A.8) \quad \begin{aligned} & \sup_{0 < t \leq \epsilon} |S_n(t)|/q(t) \\ &= n^{-1/2} \sup_{1/(n+1) \leq t \leq \epsilon} \left| \sum_{1 \leq i \leq (n+1)t} \xi_i \right| /q(t) \\ &\leq n^{-1/2} \sup_{1/(n+1) \leq t \leq \epsilon} \left| \sum_{1 \leq i \leq (n+1)t} \xi_i - \sigma_n W_n^*((n+1)t) \right| /q(t) \\ &\quad + n^{-1/2} \sup_{1/(n+1) \leq t \leq \epsilon} \sigma_n |W_n^*((n+1)t)|/q(t). \end{aligned}$$

By C.2 we have

$$\begin{aligned} & n^{-1/2} \sup_{1/(n+1) \leq t \leq \epsilon} \left| \sum_{1 \leq i \leq (n+1)t} \xi_i - \sigma_n W_n^*((n+1)t) \right| /q(t) \\ &\leq \sup_{1 \leq k \leq n} \left| \sum_{1 \leq i \leq k} \xi_i - \sigma_n W_n^*(k) \right| /k^{1/2} \sup_{0 < t \leq \epsilon} \frac{((n+1)t)^{1/2} n^{-1/2}}{q(t)} \\ &= O_P(1) \sup_{0 < t \leq \epsilon} \frac{t^{1/2}}{q(t)} \end{aligned}$$

and therefore by (A.4) we have

$$(A.9) \quad \lim_{\epsilon \rightarrow 0} \limsup_{n \rightarrow \infty} P \left\{ n^{-1/2} \sup_{1/(n+1) \leq t \leq \epsilon} \left| \sum_{1 \leq i \leq (n+1)t} \xi_i - \sigma_n W_n^*((n+1)t) \right| /q(t) > \delta \right\} = 0$$

for all  $\delta > 0$ . Observing that

$$(A.10) \quad n^{-1/2} \sup_{1/(n+1) \leq t \leq \epsilon} |W_n^*((n+1)t)|/q(t) \\ \stackrel{D}{=} \left(\frac{n}{n+1}\right)^{1/2} \sigma_n \sup_{1/(n+1) \leq t \leq \epsilon} |W(t)|/q(t),$$

we get (cf. Csörgő and Horváth (1983), p. 189) that

$$(A.11) \quad \lim_{\epsilon \rightarrow 0} \sup_{0 < t \leq \epsilon} |W(t)|/q(t) = 0 \quad \text{a.s.}$$

Putting together (A.7)–(A.11) we obtain (A.6). By the stationarity of  $\{\xi_i, -\infty < i < \infty\}$  and C.2 we get similarly to (A.6) that

$$(A.12) \quad \lim_{\epsilon \rightarrow 0} \limsup_{n \rightarrow \infty} P \left\{ \sup_{1-\epsilon \leq t < 1} |S_n^0(t)|/q(t) > \delta \right\} = 0$$

for all  $\delta > 0$ . Similarly to (A.5) we have

$$(A.13) \quad \lim_{\epsilon \rightarrow 0} \sup_{1-\epsilon < t < 1} |B(t)|/q(t) = 0 \quad \text{a.s.}$$

Now (A.2), (A.5), (A.6), (A.12) and (A.13) yield the first part of Theorem A.1.

Next we assume that (A.1) holds. First we show that  $\lim_{t \rightarrow 0} t/q(t) = 0$ . By definition,  $S_n(t) = 0$ , if  $0 \leq t < 1/(n+1)$ , and therefore by C.1 we have

$$(A.14) \quad \sup_{0 < t < 1/(n+1)} |S_n^0(t)|/q(t) = |S_n(1)| \sup_{0 < t < 1/(n+1)} t/q(t) = O_P(1)$$

and

$$(A.15) \quad \sup_{n/(n+1) < t < 1} |S_n^0(t)|/q(t) = O_P(1).$$

Hence (A.1) yields

$$(A.16) \quad \sup_{0 < t < 1/(n+1)} |B_n(t)|/q(t) = O_P(1)$$

and

$$(A.17) \quad \sup_{n/(n+1) < t < 1} |B_n(t)|/q(t) = O_P(1).$$

The distribution of  $B_n$  does not depend on  $n$ , and therefore  $I_{0,1}(q, c) < \infty$  for some  $c > 0$  (cf. Csörgő and Horváth (1993), p. 189). Since  $I_{0,1}(q, c) < \infty$  for some  $c > 0$ , we also have (A.4), and therefore (A.14) and (A.15) can be replaced by

$$(A.18) \quad \sup_{0 < t < 1/(n+1)} |S_n^0(t)|/q(t) = o_P(1)$$

and

$$(A.19) \quad \sup_{n/(n+1) < t < 1} |S_n^0(t)|/q(t) = o_P(1).$$

Thus (A.1) yields

$$(A.20) \quad \sup_{0 < t < 1/(n+1)} |B_n(t)|/q(t) = o_P(1)$$

and

$$(A.21) \quad \sup_{n/(n+1) < t < 1} |B_n(t)|/q(t) = o_P(1).$$

The distribution of  $B_n$  does not depend on  $n$ , and therefore by Csörgő and Horváth ((1993), p. 189) we conclude that  $I_{0,1}(q, c) < \infty$  for all  $c > 0$ .

The next theorem shows that the weighted supremum functionals of  $S_n^0$  converges in distribution under weaker conditions than needed for weak convergence in weighted metrics. Let  $\{B(t), 0 \leq t \leq 1\}$  be a Brownian bridge.

**THEOREM A.2.** *We assume that C.1, C.2 hold and  $q \in Q$ .  $I_{0,1}(q, c) < \infty$  for some  $c > 0$  if and only if*

$$(A.22) \quad \sup_{0 < t < 1} |S_n^0(t)|/q(t) \xrightarrow{D} \sigma \sup_{0 < t < 1} |B(t)|/q(t)$$

and

$$(A.23) \quad \sup_{0 < t < 1} S_n^0(t)/q(t) \xrightarrow{D} \sigma \sup_{0 < t < 1} B(t)/q(t).$$

**PROOF.** First we note, that if (A.22) and (A.23) hold true, then the limit distributions are almost surely finite and therefore  $I_{0,1}(q, c) < \infty$  for some  $c > 0$  (cf. Csörgő and Horváth (1993), p. 188).

Next we prove that if  $I_{0,1}(q, c) < \infty$  for some  $c > 0$ , then (A.22) also holds. Similar arguments give (A.23). Given  $q$ , by Lemma 4.1.1 in Csörgő and Horváth (1993), there are constants  $0 \leq c_1 = c_1(q) < \infty$  and  $0 \leq c_2 = c_2(q) < \infty$  such that

$$(A.24) \quad \lim_{\epsilon \rightarrow 0} \sup_{0 < t \leq \epsilon} |B(t)|/q(t) = c_1 \quad \text{a.s.},$$

$$(A.25) \quad \lim_{\epsilon \rightarrow 0} \sup_{1-\epsilon \leq t < 1} |B(t)|/q(t) = c_2 \quad \text{a.s.}$$

and

$$(A.26) \quad \lim_{\epsilon \rightarrow 0} \sup_{0 < t \leq \epsilon} |W(t)|/q(t) = c_1 \quad \text{a.s.},$$

$$(A.27) \quad \lim_{\epsilon \rightarrow 0} \sup_{0 < t \leq \epsilon} |W(t)|/q(1-t) = c_2 \quad \text{a.s.},$$

where  $\{W(t), 0 \leq t < \infty\}$  denotes a Wiener process. We use the construction in Proof of Theorem A.1. It follows from (A.2) that

$$(A.28) \quad \sup_{\epsilon \leq t \leq 1-\epsilon} |S_n^0(t) - \sigma B_n(t)|/q(t) = o_P(1)$$

$\xi_i, -\infty <$

em A.1.  
= 0. By

<  $\infty$  for  
, for some  
ced by

for all  $0 < \epsilon < 1/2$ . Next we show that

$$(A.29) \quad \lim_{\epsilon \rightarrow 0} \limsup_{n \rightarrow \infty} P \left\{ \left| \sup_{0 < t \leq \epsilon} |S_n^0(t)|/q(t) - c_1 \right| > \delta \right\} = 0$$

and

$$(A.30) \quad \lim_{\epsilon \rightarrow 0} \limsup_{n \rightarrow \infty} P \left\{ \left| \sup_{1-\epsilon \leq t < 1} |S_n^0(t)|/q(t) - c_2 \right| > \delta \right\} = 0$$

for all  $\delta > 0$ . Following the proofs in (A.7) and (A.8) we get that

$$(A.31) \quad \lim_{\epsilon \rightarrow 0} \limsup_{n \rightarrow \infty} P \left\{ \left| \sup_{0 < t \leq \epsilon} |S_n(t)|/q(t) - n^{-1/2} \sigma_n \sup_{1/(n+1) \leq t \leq \epsilon} |W_n^*((n+1)t)|/q(t) \right| > \delta \right\} = 0$$

for all  $\delta > 0$ , where  $W_n^*$  are the Wiener processes in C.2. Using (A.10) and (A.26) we get that

$$(A.32) \quad \lim_{\epsilon \rightarrow 0} \limsup_{n \rightarrow \infty} P \left\{ \left| n^{-1/2} \sigma_n \sup_{1/(n+1) \leq t \leq \epsilon} |W_n^*((n+1)t)|/q(t) - c_1 \right| > \delta \right\} = 0$$

for all  $\delta > 0$ . Now it is clear that (A.29) follows from (A.31) and (A.32).

#### REFERENCES

- Bai, J. (1994). Least squares estimation of a shift in linear processes, *J. Time Ser. Anal.*, **15**, 453–472.
- Csörgő, M. and Horváth, L. (1988). Nonparametric methods for change-point problems, *Handbook of Statistics*, **7**, 275–280.
- Csörgő, M. and Horváth, L. (1993). *Weighted Approximations in Probability and Statistics*, Wiley, New York.
- Csörgő, M. and Révész, P. (1981). *Strong Approximations in Probability and Statistics*, Academic Press, New York.
- Darling, D. A. and Erdős, P. (1956). A limit theorem for maximum of normalized sums of independent random variables, *Duke Math. J.*, **23**, 143–155.
- Davis, R. A., Huang, D. and Yao, Y. C. (1995). Testing for a change in the parameter values and order of an autoregressive model, *Ann. Statist.*, **23**, 283–304.
- Giraitis, L. and Leipus, R. (1990). A functional CLT for nonparametric estimates of spectra and change-point problem for spectral function, *Lietuvos Matematikos Rinkinys*, **30**, 674–697.
- Giraitis, L. and Leipus, R. (1992). Testing and estimating in the change-point problem of the spectral function, *Lietuvos Matematikos Rinkinys*, **32**, 20–38.
- Gombay, E. and Horváth, L. (1990). Asymptotic distributions of maximum likelihood tests for change in the mean, *Biometrika*, **77**, 411–414.
- Gombay, E. and Horváth, L. (1994). An application of the maximum likelihood test to the change-point problem, *Stochastic Process. Appl.*, **50**, 161–171.
- Gorodetskii, V. V. (1977). On the strong mixing property for linear sequences, *Theory Probab. Appl.*, **22**, 411–413.
- Haccou, P., Meelis, E. and Van de Geer, S. (1988). The likelihood ratio test for the change point problem for exponentially distributed random variables, *Stochastic Process. Appl.*, **27**, 121–139.

- Horváth, L. (1993a). Change in autoregressive processes, *Stochastic Process. Appl.*, **44**, 221–242.
- Horváth, L. (1993b). The maximum likelihood method for testing changes in the parameters of normal observations, *Ann. Statist.*, **21**, 671–680.
- Horváth, L. and Kokoszka, P. (1995). The effect of long-range dependence on change-point estimators (preprint).
- Ibragimov, I. A. and Linnik, Y. V. (1971). *Independent and Stationary Sequences of Random Variables*, Wolters-Noordhoff, Groningen.
- Kuelbs, J. and Philipp, W. (1980). Almost sure invariance principles for partial sums of mixing  $B$ -valued random variables, *Ann. Probab.*, **8**, 1003–1036.
- Kulpelger, R. J. (1985). On the residuals of autoregressive processes and polynomial regression, *Stochastic Process. Appl.*, **21**, 107–118.
- Page, E. S. (1954). Continuous inspection schemes, *Biometrika*, **41**, 100–105.
- Page, E. S. (1955). A test for change in a parameter occurring at an unknown point, *Biometrika*, **42**, 523–526.
- Philipp, W. and Stout, W. (1975). Almost sure invariance principles for partial sums of weakly dependent random variables, *Mem. Amer. Math. Soc.*, **161**, 1–137.
- Picard, D. (1985). Testing and estimating change-points in time series, *Adv. in Appl. Probab.*, **17**, 841–867.
- Sen, A. and Srivastava, M. S. (1975a). On tests for detecting change in the mean, *Ann. Statist.*, **3**, 98–108.
- Sen, A. and Srivastava, M. S. (1975b). Some one-sided tests for change in level, *Technometrics*, **17**, 61–64.
- Shorack, G. R. and Wellner, J. A. (1986). *Empirical Processes with Applications to Statistics*, Wiley, New York.
- Tang, S. M. and MacNeill, I. B. (1993). The effect of serial correlation on tests for parameter change at unknown time, *Ann. Statist.*, **21**, 552–575.
- Withers, C. S. (1981). Conditions for linear processes to be strong-mixing, *Zeit. Wahrscheinlichkeitsth.*, **57**, 477–480.
- Worsley, K. J. (1979). On the likelihood ratio test for a shift in location of normal populations, *J. Amer. Statist. Assoc.*, **74**, 365–367.
- Worsley, K. J. (1986). Confidence regions and tests for a change-point in a sequence of exponential random variables, *Biometrika*, **73**, 91–104.
- Yao, Y. C. and Davis, R. A. (1986). The asymptotic behavior of the likelihood ratio statistic for testing a shift in mean in a sequence of independent normal variates, *Sankhyā Ser. A*, **48**, 339–353.